

EconoTaq[®] PLUS 2X Master Mix

IMPORTANT!
-20°C Storage Required
Immediately Upon Receipt

Lucigen[®] Corporation
Advanced Products for Molecular Biology

2120 W. Greenview Drive
Middleton, WI 53562 USA
Toll Free: (888) 575-9695
Phone: (608) 831-9011
FAX: (608) 831-9012
E-mail: lucigen@lucigen.com
Internet: www.lucigen.com

EconoTaq[®] PLUS 2X Master Mix

Table of Contents

| | |
|--|----------|
| Components & Storage Conditions | 3 |
| Description..... | 3 |
| Introduction to PCR..... | 3 |
| PCR Setup..... | 4 |
| PCR Cycling Conditions | 4 |
| PCR Guidelines..... | 5 |
| Quality Control..... | 5 |
| Related Lucigen Products | 6 |

PLEASE NOTE

Some applications in which Lucigen's EconoTaq DNA Polymerase can be used may be covered by patents issued and applicable in the United States and certain other countries. Because purchase of this product does not include a license to perform any patented application, users of this product may be required to obtain a patent license depending upon the particular application in which the product is used. The PCR process is the subject of European Patent Nos. 201,184 and 200,262 owned by Hoffman-LaRoche. Those patents expired on March 28, 2006. The corresponding PCR process patents in the United States expired on March 29, 2005. It is the sole responsibility of the buyer to ensure that use of the product does not infringe the patent rights of third parties.

Warranty

Lucigen warrants that its products will conform to the standards stated in its product specification sheets in effect at the time of shipment. Lucigen will replace, free of charge, any product that does not conform to the specifications. This warranty limits Lucigen's liability only to the replacement of the product. THIS WARRANTY IS EXCLUSIVE, AND LUCIGEN MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING WITHOUT LIMITATION, ANY IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. The stated express warranties, and the remedy provided for breach thereof, are in lieu of all other liability or obligations of Lucigen for any damages whatsoever arising out of or in connection with the delivery, use, performance, or the inability to use any of its products. IN NO EVENT SHALL LUCIGEN BE LIABLE UNDER ANY LEGAL THEORY (INCLUDING BUT NOT LIMITED TO CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT, OR WARRANTY OF ANY KIND) FOR ANY INDIRECT, SPECIAL, INCIDENTAL, CONSEQUENTIAL, OR EXEMPLARY DAMAGES (INCLUDING BUT NOT LIMITED TO LOST PROFITS). Without limiting the effect of the preceding sentence, Lucigen's maximum liability, if any, shall not exceed the purchase price paid by buyer for the product.

Notice of Limited Label License, Copyright, Patents, Warranties, Disclaimers, and Trademarks

Copyright© 2010 by Lucigen Corp. All rights reserved. UltraClone, CloneDirect, pcrSMART, pEZ, pEZSeq, cSMART, PCR-SMART, Replicator, OverExpress, and FLASH are trademarks of Lucigen Corp. EconoTaq, BigEasy, Lucigen, CloneSmart, ClonePlex, DNATerminator, E. cloni, PCRTerminator, CopyRight, NanoClone, and pSMART are registered trademarks of Lucigen Corp.

Lucigen's products are sold for research use only and are not to be used in humans or for medical diagnostics. Lucigen's liability with respect to any of its products is limited to the replacement of the product. No other warranties of any kind, expressed or implied, including without limitation, any implied fitness for any particular use, are provided by Lucigen. Lucigen is not liable for any direct, indirect, incidental or consequential damages arising out of or in connection with the use or inability to use any of its products.

If the purchaser is not willing to accept these use limitations, Lucigen Corporation is willing to accept return of the product for a full refund. For information on obtaining a license, contact Lucigen Corporation, 2120 W. Greenview Dr., Middleton, WI 53562. Email: Lucigen@lucigen.com. Phone: 608-831-9011. Fax 608-831-9012.

EconoTaq[®] PLUS 2X Master Mix

Components & Storage Conditions

Components: EconoTaq PLUS 2X Master Mix contains all of the components necessary to perform PCR amplification, except for template and primers. The Master Mix also contains agarose gel loading buffer.

| Catalog # | Number of reactions |
|-----------|---|
| 30035-0 | 50 reactions (Trial Size: 1 tube, containing 1,250µl) |
| 30035-1 | 500 reactions (10 tubes, each containing 1,250µl) |
| 30035-2 | 1000 reactions (20 tubes, each containing 1,250µl) |

Storage: EconoTaq PLUS 2X Master Mix can be stored at –20°C for 12 months. For everyday use, an aliquot can be stored at +4°C for up to three months. The EconoTaq PLUS 2X Master Mix needs to be mixed well prior to use. It is stable for ten freeze-thaw cycles.

Description

EconoTaq PLUS 2X Master Mix is a ready-to-use PCR master mix containing agarose gel loading buffer. A PCR reaction is set up simply by combining the EconoTaq PLUS Master Mix with template DNA, primers, and water. After the reaction is complete, it can be loaded directly onto an agarose gel. All other necessary components are provided in the Master Mix, which contains the following: 0.1 units/µl of EconoTaq DNA Polymerase, Reaction Buffer (pH 9.0), 400 µM dATP, 400 µM dGTP, 400 µM dCTP, 400 µM dTTP, 3 mM MgCl₂, and a proprietary PCR Enhancer/Stabilizer. For difficult templates, such as those containing GC content up to 75%, it has been found that EconoTaq PLUS Master Mix performs well with denaturation temperatures up to 98°C.

Introduction to PCR

The Polymerase Chain Reaction (PCR) is a powerful technique that amplifies specific DNA sequences using multiple cycles of a 3-step process. The first step involves denaturation of a double-stranded DNA template at a high temperature. In the second step, sequence-specific primers anneal to complementary sites flanking the target sequence. In the third step, a thermostable DNA polymerase extends the annealed primers, thereby copying the original DNA sequence. The newly synthesized DNA becomes the template for subsequent DNA amplification, doubling the amount of template with each cycle. These steps are typically repeated 25-35 times, resulting in a 10⁵-10⁹ fold increase in the amount of target DNA.

EconoTaq[®] PLUS 2X Master Mix

PCR Setup

1) Materials supplied by the user. PCR amplification is performed by adding template DNA, primers, and water to the EconoTaq 2X Master Mix. The following components must be supplied by the user:

Template DNA (10-50 ng of plasmid DNA; 50-200 ng of genomic DNA)
Forward Primer (100 pmol/μl)
Reverse Primer (100 pmol/μl)
Nuclease-free water
Thermocycling apparatus

2) Reaction Setup. Set up PCR amplifications of the desired size, according to the following chart:

| | 25 μl Reaction | 50 μl Reaction | 100 μl Reaction | Final concentration |
|------------------------------|-------------------|-------------------|--------------------|------------------------|
| EconoTaq PLUS 2X Master Mix | 12.5 μl | 25.0 μl | 50.0 μl | 1 X |
| Forward Primer (100 pmol/μl) | 0.25 μl | 0.5 μl | 1.0 μl | 1 pmol/μl (1 μM) |
| Reverse Primer (100 pmol/μl) | 0.25 μl | 0.5 μl | 1.0 μl | 1 pmol/μl (1 μM) |
| DNA template (10 ng/μl) | 1.0 μl | 1.0 μl | 1.0 μl | |
| Water, Nuclease-free | 11.0 μl | 23.0 μl | 47.0 μl | |

3) Gently mix the PCR components in a thin-walled reaction tube and spin briefly in a microcentrifuge. Add a drop of mineral oil if the thermocycler does not have a heated lid.

PCR Cycling Conditions

- 1) Pre-heat the thermocycler to 94°C.
- 2) For initial denaturation of target template DNA, incubate the reactions at 94°C for two minutes.
- 3) Denature, anneal, and extend the DNA according to the following chart for subsequent cycles of amplification:

| Cycling step | Temperature | Time | # of Cycles |
|-----------------------|-------------|--------------|-------------|
| Initial Denaturation* | 94°C | 2 min | 1 |
| Denaturation* | 94°C | 15 - 30 sec | 25 - 35 |
| Annealing** | 50 - 65°C | 15 - 30 sec | |
| Extension | 72°C | 1 min/kb | |
| Final Extension | 72°C | 5 - 10 min | 1 |
| Hold | 4°C | Indefinitely | 1 |

* EconoTaq PLUS performs well with denaturation temperatures up to 98°C.

** Anneal at T_m of primer $\pm 2^\circ\text{C}$. See PCR Guidelines section.

4) After completion of the PCR, a 5 μL aliquot of the reaction is loaded onto an agarose gel for analysis or size selection.

EconoTaq[®] PLUS 2X Master Mix

PCR Guidelines

Optimization of PCR conditions is required for amplification involving templates with high GC content, internal secondary structure, or products greater than 5 kb. The following guidelines can be used to improve the success of amplifying these templates:

- 1) **Cold Reaction Set-Up.** Reactions using EconoTaq PLUS Master Mix yield best results when set up on ice and maintained at 4°C prior to amplification. The EconoTaq Polymerase has residual activity at temperatures above 4°C that can cause non-specific background amplification. Add primers last to reactions just prior to target addition and incubation/cycling.
- 2) **Template DNA.** Use of purified, high quality template DNA enhances the success rate of PCR. We recommend using 10 - 50 ng of plasmid DNA and 50 - 200 ng of genomic DNA. Template should be suspended in water, rather than EDTA-containing solution such as TE buffer.
- 3) **Primer Design.** Oligonucleotide primers for PCR should be 20-25 bases in length and should have a GC content of 40-60%, with the GC bases evenly spaced in the primer. Self-annealing of primers leads to production of primer-dimers, which can diminish the amount of authentic product; therefore, the 3' end of each primer should not be complementary to itself or to the 3' end of the opposing primer. The melting temperature (T_m) of the primers should be within 5°C of each other. The final primer concentration in the reaction should be 1 pmol/ μ l (equal to 1 nmol/ml, or 1 μ M).
- 4) **Annealing.** PCR enhancers in EconoTaq PLUS Master Mix may lower primer T_m slightly. Optimize the reaction conditions by performing the reaction starting with an annealing temperature 2°C below the calculated melting temperature of the primers. Vary the annealing temperature to optimize for the primer/template combination. Reducing the annealing temperature improves yield and reaction efficiency, while increasing annealing temperature improves specificity.
- 5) **Denaturation Temperature.** EconoTaq PLUS 2X Master Mix will amplify most routine DNA templates using a denaturation temperature of 94°C. For GC-rich templates, with GC content up to 75%, a denaturation temperature of 98°C has been found to improve results.

Quality Control

PCR Activity: EconoTaq 2X Master mix is tested in DNA amplification using a variety of templates and primers.

Activity Determination: One unit of EconoTaq DNA Polymerase catalyzes the incorporation of 10 nmoles of dNTP into acid-insoluble material in 30 minutes at 70°C in 50 mM Tris-HCl (pH 9.0), 50 mM NaCl, 5 mM MgCl₂, 200 μ M dGTP, dATP, dTTP, dCTP (a mix of unlabeled and [³³P]dCTP), 10 μ g Activated Calf Thymus DNA, and 0.1 mg/ml BSA.

Absence of Endonuclease or Nicking Activity: Incubation of 10 U of EconoTaq DNA Polymerase with 1 μ g of supercoiled pBR322 DNA for 16 hours at 70°C results in no detectable conversion to relaxed or linear forms detectable by agarose gel electrophoresis.

Absence of Exonuclease Activity: Incubation of 10 U of EconoTaq DNA Polymerase with 1 μ g of HindIII-cut lambda DNA for 16 hours at 70°C resulted in no smearing of bands on agarose gels.

Purity: EconoTaq DNA Polymerase is >99% pure as determined by SDS-PAGE. There is no detectable DNA contamination.

EconoTaq[®] PLUS 2X Master Mix

Related Lucigen Products

- EconoTaq[®] PLUS GREEN 2X Master Mix
- EconoTaq DNA Polymerase
- TaqSelect[™] DNA Polymerase
- Bst DNA Polymerase, Exonuclease Minus
- PyroPhage[®] 3173 DNA Polymerase, Exonuclease Minus (Exo-)
- dNTPs, PCR Grade
- Gel-Ready[™] DNA Ladders
- BigEasy[®] v2.0 Linear Cloning Kit
- CloneSmart[®] Blunt Cloning Kits
- DNATerminator[®] End Repair Kit
- PCRTerminator[®] End Repair Kit
- UltraClone[™] DNA Ligation & Transformation Kits
- CloneDirect[™] Rapid Ligation Kit
- GC Cloning and Amplification Kits
- ClonePlex[®] Library Construction Kit
- pEZSeq[™] Blunt Cloning Kits
- cSMART[™] cDNA Cloning Kits
- *E. coli*[®] EXPRESS Competent Cells
- OverExpress[™] Competent Cells